

07-26-00

A

**UTILITY PATENT APPLICATION TRANSMITTAL**  
**(Small Entity)**

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.  
PENN-0708

bax se 2

Total Pages in this Submission

**TO THE ASSISTANT COMMISSIONER FOR PATENTS****Box Patent Application**  
**Washington, D.C. 20231**

Transmitted herewith for filing under 35 U.S.C. 111(a) and 37 C.F.R. 1.53(b) is a new utility patent application for an invention entitled:

**A RAPID, SENSITIVE AND QUANTITATIVE METHOD FOR IMMUNO-DETECTION OF EPITOPES ON MOLECULES USING A SINGLE CHAIN FV FOR THE EPITOPE OR A CONSTRAINED EPITOPE SPECIFIC CDR**

and invented by:

**Greene et al.**If a **CONTINUATION APPLICATION**, check appropriate box and supply the requisite information:☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: \_\_\_\_\_

Which is a:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: \_\_\_\_\_

Which is a:

☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: \_\_\_\_\_

Enclosed are:

**Application Elements**

1. ☒ Filing fee as calculated and transmitted as described below
2. ☒ Specification having 22 pages and including the following:
  - a. ☒ Descriptive Title of the Invention
  - b. ☐ Cross References to Related Applications (if applicable)
  - c. ☐ Statement Regarding Federally-sponsored Research/Development (if applicable)
  - d. ☐ Reference to Microfiche Appendix (if applicable)
  - e. ☒ Background of the Invention
  - f. ☒ Brief Summary of the Invention
  - g. ☐ Brief Description of the Drawings (if drawings filed)
  - h. ☒ Detailed Description
  - i. ☒ Claim(s) as Classified Below
  - j. ☒ Abstract of the Disclosure

# UTILITY PATENT APPLICATION TRANSMITTAL (Small Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.  
PENN-0708

Total Pages in this Submission

## Application Elements (Continued)

3. ☐ Drawing(s) (when necessary as prescribed by 35 USC 113)
- a. ☐ Formal      b. ☐ Informal      Number of Sheets \_\_\_\_\_
4. ☒ Oath or Declaration
- a. ☐ Newly executed (original or copy)      ☒ Unexecuted
- b. ☐ Copy from a prior application (37 CFR 1.63(d)) (for continuation/divisional application only)
- c. ☒ With Power of Attorney      ☐ Without Power of Attorney
- d. ☐ DELETION OF INVENTOR(S)  
Signed statement attached deleting inventor(s) named in the prior application,  
see 37 C.F.R. 1.63(d)(2) and 1.33(b).
5. ☐ Incorporation By Reference (usable if Box 4b is checked)  
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.
6. ☐ Computer Program in Microfiche
7. ☒ Genetic Sequence Submission (if applicable, all must be included)
- a. ☒ Paper Copy
- b. ☒ Computer Readable Copy
- c. ☒ Statement Verifying Identical Paper and Computer Readable Copy

## Accompanying Application Parts

8. ☐ Assignment Papers (cover sheet & documents)
9. ☐ 37 CFR 3.73(b) Statement (when there is an assignee)
10. ☐ English Translation Document (if applicable)
11. ☐ Information Disclosure Statement/PTO-1449      ☐ Copies of IDS Citations
12. ☐ Preliminary Amendment
13. ☒ Acknowledgment postcard
14. ☒ Certificate of Mailing
- ☐ First Class      ☒ Express Mail (Specify Label No.): EL603801965US

# UTILITY PATENT APPLICATION TRANSMITTAL (Small Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.  
PENN-0708

Total Pages in this Submission

## Accompanying Application Parts (Continued)

15. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
16. ☒ Small Entity Statement(s) - Specify Number of Statements Submitted: 1
17. ☐ Additional Enclosures (please identify below):


## Fee Calculation and Transmittal

### CLAIMS AS FILED

For	#Filed	#Allowed	#Extra	Rate	Fee
Total Claims	11	- 20 =	0	x \$9.00	\$0.00
Indep. Claims	3	- 3 =	0	x \$39.00	\$0.00
Multiple Dependent Claims (check if applicable) <input type="checkbox"/>					\$0.00
BASIC FEE					\$345.00
OTHER FEE (specify purpose)					\$0.00
TOTAL FILING FEE					\$345.00

- ☒ A check in the amount of **\$345.00** to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge and credit Deposit Account No. **12-1086** as described below. A duplicate copy of this sheet is enclosed.
- ☐ Charge the amount of \_\_\_\_\_ as filing fee.
- ☒ Credit any overpayment.
- ☒ Charge any additional filing fees required under 37 C.F.R. 1.16 and 1.17.
- ☐ Charge the issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

Dated: July 25, 2000



Signature

Jane Massey Licata  
Reg. No. 32,257  
Law Offices of Jane Massey Licata  
66 E. Main Street  
Marlton, NJ 08053  
Tel: 856-810-1515  
Fax: 856-810-1454

CC:

PENN-0708

CERTIFICATE OF EXPRESS MAILING

"Express Mail" Label No.EL603801965US

Date of Deposit JULY 25, 2000

I hereby certify that this paper is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the "BOX SEQUENCE", Assistant Commissioner for Patents, Washington, D.C. 20231.

- 1) Patent Application Transmittal Letter (2 copies);
- 2) Application consisting of 22 pages of Specification, including two (2) pages of Claims, and one (1) page of Abstract;
- 3) Return Post Card;
- 4) Check in the amount of \$345.00;
- 5) Verified Statement Claiming Small Entity Status;
- 6) Unexecuted Declaration and Power of Attorney;
- 7) Statement to Support Filing and Submission in Accordance with 37 CFR §§1.821-1.825;
- 8) Sequence listing; and
- 9) Diskette containing computer readable copy of Sequence Listing.

Jane Massey Licata  
JANE MASSEY LICATA



009220 94649460



Each person, concern or organization to which I have assigned, granted, conveyed, or licensed or am under an obligation under contract or law to assign, grant, convey, or license any rights in the invention is listed below:

- ☒ no such person, concern or organization exists.  
☐ each such person, concern or organization is listed below.

FULL NAME \_\_\_\_\_  
ADDRESS \_\_\_\_\_

☐ Individual ☐ Small Business Concern ☐ Nonprofit Organization

FULL NAME \_\_\_\_\_  
ADDRESS \_\_\_\_\_

☐ Individual ☐ Small Business Concern ☐ Nonprofit Organization

FULL NAME \_\_\_\_\_  
ADDRESS \_\_\_\_\_

☐ Individual ☐ Small Business Concern ☐ Nonprofit Organization

FULL NAME \_\_\_\_\_  
ADDRESS \_\_\_\_\_

☐ Individual ☐ Small Business Concern ☐ Nonprofit Organization

Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING: Alan Dickason  
TITLE IN ORGANIZATION: Director,  
ADDRESS OF PERSON SIGNING: Center for Technology Transfer  
3700 Market Street, Suite 300  
Philadelphia, PA 19104

SIGNATURE: Alan Dickason DATE: 7/25/00  
Director, Learning & Sciences

A RAPID, SENSITIVE AND QUANTITATIVE METHOD FOR  
IMMUNO-DETECTION OF EPITOPES ON MOLECULES  
USING A SINGLE CHAIN FV FOR THE EPITOPE OR A CONSTRAINED  
EPITOPE SPECIFIC CDR

5 Background of the Invention

One of the central problems in cell biology and medicine relates to the inability to monitor protein, lipids, sugars and metabolite levels and their modifications in the single living cell. A variety of  
10 technologies have been employed to improve the sensitivity of detecting these molecules.

For example, to increase the sensitivity of immunoassays able to detect proteins at very low amounts, the polymerase chain reaction (PCR) technology has been  
15 combined with conventional immuno-detection methods ((U.S. Patent 5,665,539). This technology, termed immuno-PCR, provides an extremely sensitive method to detect proteins. In immuno-PCR, a linker molecule with bi-specific binding affinity for DNA and antibody is used to attach a marker  
20 DNA molecule specifically to an antigen-antibody complex, thus resulting in the formation of a specific antigen-antibody-DNA conjugate. The attached marker DNA can be amplified by PCR with the appropriate primers. As described in U.S. Patent 5,665,539, antigen is immobilized on the  
25 surface of microtiter plates and subsequently detected by immuno-PCR. Using this technique, an approximately  $10^5$  increase in sensitivity over an alkaline phosphatase conjugated ELISA was obtained. Sensitivity advantages of immuno-PCR have subsequently been confirmed in assays for  
30 mouse anti-lipoprotein IgG (Ruzicka et al. Science 1993

260:698-699); a human proto-oncogene protein (Zhou et al. Nucleic Acid Res. 1993 21:6038-6039); and tumor necrosis factor alpha (Sanna et al. Proc. Natl. Acad. Sci. 1995 92:272-275).

5           However, the original immuno-PCR protocol used a streptavidin-protein A chimera to detect the antigen-antibody complex. The variation in the affinity of protein A against different classes of IgGs limits its direct application in the detection of a broad range of  
10           antigens. Certain improved protocols tried to solve this problem by introducing biotinylated secondary antibody or free streptavidins.

          Joerger et al. (Clin. Chem. 1995 41(9): 1371-1377) demonstrated that double-stranded DNA labels can be  
15           directly attached to antibodies, thus allowing conjugate reagents to be prepared before the assay.

          Suzuki et al. (Jpn. J. Cancer Res. 1995 86:885-88) describe a method called double determinant  
immuno-polymerase chain reaction (double-determinant  
20           immuno-PCR) which utilizes two monoclonal antibodies, in which the antigens are sandwiched, and a specific DNA molecule is used as a marker. In this method, the first monoclonal antibody to bind the circulating antigen is immobilized instead of the antigen itself. A biotinylated  
25           second monoclonal antibody is bound to the antigen and free streptavidin is used to attach a biotinylated DNA to the second monoclonal antibody. The biotinylated DNA complexed with antigen-antibody-streptavidin is amplified by PCR. The products are then analyzed by Southern blot analysis.

30           While these immuno-PCR techniques have provided advantages over traditional methods of protein detection such as an increase in sensitivity, there still exist several notable limitations to their use. One of the major limitations of immuno-PCR lies in the non-linear  
35           amplification ability of PCR reaction. There is no direct



correlation between the amount of signal and the amount of protein present. Thus, this technique is limited as a quantitative detection method.

U.S. Patent 5,922,553 discloses a method for  
5 quantifying levels of a selected protein via a technique referred to as immuno-aRNA. In this method, a first antibody targeted to a selected protein is immobilized to a solid support. The support is then contacted with the selected protein so that the selected protein is  
10 immobilized to the first antibody. The solid support is then contacted with a RNA promoter-driven cDNA sequence covalently coupled to a second antibody targeted to the selected protein so that the second antibody binds to the bound selected protein. The amount of selected protein is  
15 determined by quantifying levels of the promoter driven cDNA sequence covalently coupled to the bound second antibody via an amplified RNA technique. In a preferred embodiment, a T7 promoter driven cDNA sequence is covalently coupled to the second antibody.

20 It has now been found that single chain fragments as well as exocyclic peptide based complementarity determining region (CDR) subunits can be used in this immuno-aRNA technique. Further, it has been found that PCR, as well as amplified RNA techniques, can be used to quantify the  
25 promoter driven cDNA sequence covalently coupled to the bound single chain fragment or CDR subunit. The use of smaller antibody binding units and fragments coupled with the already existing large single chain or cyclic peptide libraries and the use of robotic assistance renders this  
30 method widely useful for both medicinal and research purposes. Furthermore, a single third detector species can be coupled with double-stranded DNA and bound to either the single chain Fv or the CDRs, rendering detection uniform and simple. This is referred to herein as a universal  
35 detector.

Summary of the Invention

An object of the present invention is to provide a method for detecting molecules expressing a selected epitope in a sample. In this method, an epitope anchor  
5 specific for a selected epitope is immobilized to a selected surface. The epitope anchor may comprise a single chain Fv fragment, a CDR, an antibody, or other ligand peptide or chemical or pharmaceutical that interacts with a selected epitope. The surface is then contacted with a  
10 sample suspected of containing molecules which express the selected epitope so that the molecules bind to the immobilized epitope anchor. An epitope detector comprising a single chain Fv for the selected epitope or a constrained epitope specific CDR attached to an oligonucleotide is then  
15 used to detect any bound molecules. In one embodiment, the single chain Fv or the CDR has been modified to allow for attachment of oligonucleotides to a single site. Alternatively, the method of the present invention can be performed with an epitope anchor. In this embodiment, the  
20 epitope detector is employed to define molecules bound directly to a surface.

Another object of the present invention is to provide systems for the detection of molecules expressing a selected epitope. These systems of the present invention  
25 comprise an epitope anchor specific for a selected epitope, a selected surface on which the epitope is or can be immobilized, and an epitope detector comprising a single chain Fv for the selected epitope or a constrained epitope specific CDR attached to an oligonucleotide. In one  
30 embodiment, the single chain Fv or the constrained epitope specific CDR is modified to allow for attachment of the oligonucleotides.

Yet another object of the present invention is to provide kits for the detection of molecules expressing a  
35 selected epitope which comprise an epitope anchor specific

for a selected epitope and an epitope detector comprising a single chain Fv for the selected epitope or a constrained epitope specific CDR attached to an oligonucleotide. In one embodiment, the single chain Fv or the constrained  
5 epitope specific CDR is modified to allow for attachment of the oligonucleotides.

### Detailed Description of the Invention

The present invention relates to improved methods for quantifying levels of a selected molecule and systems and  
10 kits for performing these improved methods. In one embodiment, the method comprises binding an epitope anchor specific for a selected epitope of the molecule to a selected surface. The epitope anchor may comprise a single chain Fv fragment, a CDR, an antibody, or other ligand  
15 peptide or chemical specific for a selected epitope. In a preferred embodiment, the epitope anchor is bound to a designated spot on the surface. For example, the surface may comprise a chip and the epitope anchor is bound to a defined spot on the chip. In one embodiment, the epitope  
20 anchor is deposited onto a surface or plate with the aid of a pipettor or similar device which permits application at a single site. The surface with the bound epitope anchor is then contacted with a sample suspected of containing molecules expressing the selected epitope so that the  
25 molecule binds to the epitope anchor. In another embodiment, the molecule is attached to a surface directly, without the use of an epitope anchor.

Examples of samples which can be assayed via the methods of the present invention include, but are not  
30 limited to, individual cells and solutions including biological fluids such as serum. An epitope detector which can bind to any bound molecule on the surface is then used to detect and quantify the amount of bound molecule. The epitope detector comprises a single chain Fv for the

selected epitope or a constrained epitope specific CDR which have been modified to allow for attachment of oligonucleotides at a single site.

Fv fragments for selected epitopes can be produced in  
5 cells or on microorganisms by use of recombinant DNA technology. For example, Skerra and Pluckthun (Science 1988 240:1038-1044) describe an expression system for production of functional Fv fragments in *E. coli*.

A method for producing Fv fragments in eukaryotic  
10 host cells with a eukaryotic expression vector which has an operon having a DNA sequence which encodes the variable domain only of an antibody light or heavy chain has also been described (J. Mol. Biol. 1988 203:825-828). Chains of the Fv fragment are secreted and correctly assembled by the  
15 host cell such that fully functional Fv fragments are produced in the culture supernatant. In addition, the DNA coding sequence may be altered toward its 5' end so that the amino terminal end expresses a residue or residues with a surface suitable for covalent coupling of an  
20 oligonucleotide. In addition, the 3' terminal end may be varied so that cysteine residues are produced towards the C-terminal end of each variable domain permitting the variable domains in the dimer to become linked together by disulphide bonding. This may also promote assembly of the  
25 Fv fragment. Alternatively, the Fv fragment may be stabilized by use of a vector having a first DNA sequence encoding a first variable domain and a second DNA sequence encoding a second variable domain, the first and second sequences being linked by a third DNA sequence which  
30 encodes a joining peptide sequence. In this case, the joining peptide sequence is sufficiently long and flexible to allow folding of the two polypeptides into a functional single chain Fv. Preferably, the host cell is a myeloma cell line which, prior to transformation, does not secrete  
35 whole antibody or light chains. Such cells lines are well

known and widely available (Reichmann et al. J. Mol. Biol. 1988 203:825-828).

It is believed that random phage technology to any hapten or chemical compound can also be used to select Fvs.  
5 (Harrison et al. United States Biochemical Pharma Ltd. (Europe), Watford, United Kingdom)

The CDR technology is well known and has been described in U.S. Patent 5,334,702, U.S. Patent 5,663,144, and U.S. Patent 5,919,764. In general, CDRs comprises a 6  
10 to 15 mer peptide constrained to be cyclic and modified by aromatic residues. An important step in the design of conformationally constrained peptides for use in the present invention is the delineation of the residues that are important for activity. This is generally accomplished  
15 by first synthesizing a set of analogs from the bioactive domain of the original antibody or receptor or ligand of different lengths and establishing the minimal chain lengths for the complete and partial activities. Once the minimal chain length has been established, each side chain  
20 can be systematically varied to determine the importance of charge, steric bulk, hydrophobicity, aromaticity, and chirality at each position. After evaluation of the properties of a large set of analogs, it is possible to identify the functional groups and conformation features  
25 involved in binding. Different conformationally constrained analogs can then be developed. Various means for constraining peptides have been developed.

One means involves introducing a conformationally constrained amino acid. Hruby (Life Sci. 1982 31:189-199)  
30 describes the synthesis of a large number of amino acid and dipeptide derivatives with built-in conformational constraints, as well as their incorporation into biologically active peptides. Prasad et al. (Biopolymers 1995 35:11-20) also describes a method of constraining the  
35 conformation of an amino acid unit by replacing the

hydrogen atom at the  $\alpha$ -carbon with a methyl group to produce a dialkylamino acid. U.S. Patent 6,022,523 describes a method that restricts the conformational freedom of amino acids by introducing a double-bond at the C- $\alpha$  and C- $\beta$  atoms.

Another means for constraining peptides involves introduction of covalent cross-links. Constraining the peptide backbone by introduction of covalent cross-links provides more dramatic effects than incorporating unusual amino acids. Macrocyclization is often accomplished by forming an amide bond between the peptide N- and C-termini, between a side chain and the N or C terminus, or between two side chains. A head-to-tail cyclization of side protected peptides synthesized by Fmoc/t-butyl solid phase procedures on polystyrene resin derivatized with 4-hydroxymethyl-3-methoxyphenoxyacetic acid, the first generation dialkoxy-benzyl linkage agent, has been described by Sheppard, R.C. (Int. J. Peptide Res. 1982 20:451-454). In addition, the analogous linkage agent, 4-(4-hydroxymethyl-3-methoxyphenoxy)-butyric acid (HAMA), was recently employed in fragment condensation and solid phase synthesis of peptides with these highly acid sensitive linkers (In Peptides, E. Giralt and D. Andreu eds, ESCOM, Leiden, The Netherlands 1991, 131-133). The enkephalin analogs described by Schiller provide an example of side-chain to backbone covalent cyclization in which covalent attachment of the  $\epsilon$ -amino group of the D-lys residue to the C terminal backbone carboxylate group of Leu produces a cyclic 16-membered ring analog with high potency and significant  $\mu$  receptor selectivity (Schiller et al. Int. J. Pep. Prot. Res. 1985; 25:171-177). BOP-reagent and carboimide/1-hydroxy-benzotriazole combinations have also been reported to be useful in the formation of cyclic peptides (Felix, A.M. Int. J. Pep. Prot. Res. 1988 31:231-238). Degrado et al. have also developed a biologically

active cyclized peptide analog of the GP IIb/IIIa complex using m-aminomethylbenzoic acid as the linker (U.S. Patent 6,022,523).

Disulphides can also be formed by oxidation via  
5 introduction of cysteine at certain positions. For example, Romani, S. (Int. J. Pep. Prot. Res. 1987 29:107-117) demonstrated that non-symmetrical disulphides can be built with the help of the di-tertbutyl ester of azodicarboxylic acid. Ploux, O. (Int. J. Pep. Prot. Res.  
10 1987 29:162-169) also describes a method for formation of non-symmetrical disulphides via thiol displacement of the 3S-3-nitro-2-pyridinesulphenyl group.

In a preferred embodiment, the oligonucleotide comprises a T7 promoter driven cDNA sequence so that it can  
15 be amplified using T7 RNA polymerase. In this embodiment, double stranded cDNA is synthesized for use as a template for T7 RNA polymerase transcription. T7 RNA polymerase requires its promoter site to be double stranded.

In one embodiment, the site on the Fv or CDR to which  
20 the oligonucleotides are attached comprises a series of residues which allow the attachment of linkers consisting of chemicals such as heterodimeric coupling reagents or other linkers. These residues provide a uniform binding site for the linker attachment. The linkers attach to this  
25 site and also links oligonucleotides to the Fv or CDR. Oligonucleotides may be unmodified or modified. For example, the presence of the amplified oligonucleotide can be enhanced by incorporating a beacon or fluorescent labeled oligonucleotide into the mixture allowing for rapid  
30 semi quantitative assessment of the epitope expressing molecules (Ton et al. Chemistry 2000 6:1107-1111; Leone et al. Nucleic Acids Res. 1998 26(9):2150-2155).

Bound epitope detectors may be quantified by methods such as amplification by conventional PCR or aRNA  
35 techniques. If the detection method used is immuno aRNA,

double-stranded oligonucleotides are used in the epitope detector. In this embodiment, aRNA is transcribed on the solid support using a polymerase which recognizes a specific promoter such as T7 RNA polymerase, T3 RNA  
5 polymerase or SP6 RNA polymerase, unlabeled ribonucleotides, and fluorescently labeled ribonucleotides.

A variety of means are available for detection of amplified products of the epitope detector. In one embodiment, the aRNA is detectably labeled such as with a  
10 radioactive label or a fluorescent label. Alternatively, the aRNA can be converted to cDNA and detected via methods such as gel electrophoresis, high performance liquid chromatography, hybridization assays, immunohistochemical assays and/or specific binding protein assays.

15 Use of Fvs and CDR peptides as the epitope detector renders this method useful in identifying larger polypeptides than can be identified by the immuno-aRNA technique of U.S. Patent 5,922,553, as well as in identifying motifs in supernatants, fluids, extracts of  
20 cells or bacteria or any other eukaryotic organism. Accordingly, the method of the present invention has widespread applicability in both medicinal and research purposes. Further, the method of the present invention is more sensitive than currently available methods and  
25 provides quantitative results.

The ability of Fvs and CDR peptides to serve as epitope detectors of selected molecules in the method of the present invention was demonstrated for the p185 receptor. Using this method, the released p185 receptor  
30 was detected at a  $10^8$ -fold increase in sensitivity over the ELISA and about a 1000-fold increase over the Western-ECL method. For these experiments, a single chain Fv (ScFv) construct of 7.16.4 (designated as 7.16.4 ScFv) was produced in accordance with the procedure outlined by  
35 Peterson & Greene (DNA and Cell Biology 1998 17(12):1031-



1040) wherein the Fv region of the heavy chain and light chain region was joined by a (gly4-Ser)5 linker. Since ScFv7.16.4 contained a poly-histidine tag, it was purified over Ni-NTA resin. After purification, the binding of 7.16.4ScFv was confirmed by FACS analysis on B104-1-1 cells, in both direct binding and competitive binding against the monoclonal antibody 7.16.4.

AHNP, a constrained exocyclic peptide designed from the CDR3.H region of the anti-human p185 antibody 4D5 was also used. AHNP binds to p185 and mimics the growth-inhibitory effects of 4D5 (Park et al. Nature Biotechnology 2000 18:194-198).

Both ScFv7.16.4 and AHNP were coupled to a double-stranded oligonucleotide (ds-oligo) to form epitope detectors. Both ds-oligo coupled 7.16.4ScFv and AHNP were able to detect their antigens, rat p185neu from B104-1-1 cells and human p185her2/neu from T6-17 cells, respectively. Further, conjugation with the ds-oligo to form the epitope detector did not change the binding affinity of the CDR detection molecules with their antigens as determined by plasmon resonance analysis. Since 7.16.4ScFv and mAb 7.16.4 have comparable binding affinity against the p185 receptor, they were used at comparable molar concentration in this assay. However, ANHP was used at a higher concentration since its affinity is lower against p185Her2/neu than 4D5.

Coupling of a ds-oligo directly to the CDR or single chain Fv can be a time-consuming procedure, particularly if the purpose is to detect hundreds or thousands of antigens in a mass screening proteomic assay. In addition, variation in the coupling efficiency can complicate the interpretation of the amplification results. Accordingly, in a preferred embodiment of the present invention, the Fv or CDR contains a general or universal epitope such as an hemagglutinin HA tag or polyhistidine tag. An example of a

general or universal epitope is the poly-His-tag in the 7.16.4ScFv initially designed for the purification of the protein. A single monoclonal antibody or single chain Fv coupled with ds-DNA is then used to bind to the general  
5 epitope to create a universal epitope detector. The efficacy of a universal epitope detector in the method of the present invention was demonstrated with the poly-His-tag in the 7.16.4ScFv. In these experiments, p185 receptors were captured by A11 coated on the plate, free  
10 7.16.4ScFv was added, followed by extensive washing and then incubating with ds-oligo conjugated anti-His antibody. After T7 polymerase amplification, specific bands from lysates of  $10^{-6}$  dilution of the cells were detected. Accordingly, this sensitivity was even higher than that  
15 seen with the basic protocol without a universal epitope detector.

The method of the present invention is also useful in the detection of post-translation modifications. PCR and aRNA techniques were originally developed to detect the  
20 activity of target genes at the DNA level. These methods have been adopted exclusively in the application of genomics research, sometimes combined with hybridization. Regardless of sensitivity, these methods are not able to detect the post-translation modification at the protein  
25 level. Monitoring of such events, however, is very critical since many modifications including, but not limited to, phosphorylation and glycosylation are related to the functional status of the protein. Thus, experiments were performed to demonstrate the ability of the method of  
30 the present invention to detect the phosphorylation of the p185 receptor induced by EGF treatment. A signaling model was established in which, upon EGF stimulation, EGFR heterodimerizes with and trans-activates p185, resulting in the phosphorylation of tyrosine residues on the p185  
35 receptor (Qian et al. Proc. Natl Acad. Sci. 1994 15:1500-

1504). The A431 cell line, which overexpresses EGFR as well as p185 erbB2, was used in these experiments. After EGF stimulation, the p185 receptor in the cell lysate was captured by 1E1, a monoclonal antibody developed against p185erbB2/neu. PY99, an IgG2b type of anti-phosphorylated Tyr antibody, was used to detect phosphorylated receptors. A second antibody, anti-IgG2b, coupled with ds-oligo, was used to probe the antigen-antibody sandwich complex. A431 cells stimulated with EGF produced a positive band, which was not observed in cells without EGF treatment. T6-17 cells, however, also showed a positive band, indicating constitutive phospho-tyrosine on the p185 receptor. These data indicate that this method is capable of detecting the functional status of a protein by analyzing its modification. Epitope detectors comprising an Fv or CDR coupled to the ds-oligo can also be used to detect the functional status of the protein.

Thus, the present invention provides a sensitive detection method which eliminates concerns about the non-quantitative nature of immuno-PCR techniques and which offers vast potential in the field of proteomics. By using a polymerase which recognizes a specific promoter such as T7 RNA polymerase, T3 RNA polymerase or SP6 RNA polymerase as well as the specific promoter in the amplification step, assays performed in accordance with this method possess linear amplification and precise quantification which are relevant to biological and medical assays. The number of factors that affect the sensitivity of detection have also been reduced. The specific binding between antigens and their Fv or monovalent CDR is the only critical parameter of this method.

The ability to provide universal epitope detectors provides the method of the present invention with multiple additional advantages. First, any cellular antigens can be detected without having been first coupled to a monoclonal

antibody with ds-oligo. Without the universal probe, the method would only be useful in looking at one or several particular antigens at a time. The universal probe, on the other hand, allows for the detection of any cellular or fluid residing antigen with available Fvs or CDRs. In addition, with slight modification in the protocol, different proteins can be detected simultaneously in a single electrophoresis lane when oligonucleotides of different sizes are attached to the Fv or CDR of the epitope detector. Thus, as demonstrated herein, the method of the present invention provides a versatile technique that is applicable in the identification of protein antigens as well as post-translational modification of polypeptides and other structures such as sugars or lipids at the single cell level of detection.

It is believed that this method will also be useful in the analysis of protein interactions and the detection of small molecules. For example, ligand peptides can be used as epitope detectors on tissue samples to identify the expression of specific receptors, or verse visa. With available Fvs, CDRs, or binding proteins, small molecules such as toxins or drug metabolites, can be detected in any solution including, but not limited to, water, foods, and body fluids.

The original immuno-PCR used pure antigens in the assay. Later iterations of Immuno-PCR examined mixed antigens (Hendrickson et al. Nucleic Acids Research 1999 23(3):522-529) but only showed sensitivity of two to three orders of magnitude higher than ELISA. In a real-world assay with the background comprising a huge variety of non-specific antigens, sensitivity is always limited by the specificity of the assay. Epitopes bound by the Fvs or CDR fragments are expected to identify larger polypeptides and can be used to identify motifs in supernatants, fluids, extracts of cells or bacteria or any other eukaryotic

organism. Further, actual identity of the polypeptides, organic molecules or sugar structures can be determined by computer aided analysis of data bases using the binding of several epitopes by Fvs as a guide. For example, binding  
5 by Fv a, d, e, and f would identify a sugar molecule as having side chains a, d, e, and f, and hence belonging to a family of sugars having these same side chains. In this way the present invention allows definition and identification of many, if not all molecules in a cell at  
10 any one particular time. Moreover this approach can be used to identify alternative transcriptional forms translated in an active cell or cellular supernatant. This procedure is easily amenable to 1) use with nonradioactive detection methods, 2) microtized liquid handling  
15 procedures, 3) low sample volume detection such as "protein chip" analysis and 4) robotization.

The following nonlimiting examples are provided to further illustrate the present invention.

#### EXAMPLES

##### 20 Example 1: Materials

Antibodies used include mAbs 7.16.4 and A11, each of which is reactive with a different epitope in the extracellular domain of p185neu. The polyclonal antiserum,  $\alpha$ -Bacneu, directed against the intracellular domain of  
25 p185neu, was used for Western Blotting. 1E1 is an IgG1 monoclonal antibody generated against the ectodomain of human p185her2/neu. rhuMAb 4D5 (Herceptin) was obtained from Genentech. The anti-phosphotyrosine monoclonal antibody PY99 was obtained from Santa Cruz Biotechnology  
30 (Santa Cruz, CA). The cell line B104-1-1 was derived from NIH3T3 cell by expressing rat oncogenic p185neu. The expression level of p185neu in B104-1-1 cells was determined using an  $^{125}\text{I}$ -labeled anti-neu mAb binding assay. NR6, negative for both EGFR and p185neu, was a subclone

from NIH3T3. T6-17 was derived from NIH3T3 by over-expressing the human p185her2/neu receptor. These cell lines were all cultured in Dulbecco's Modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS, 5 Hyclone) at 37°C in a 5% CO<sub>2</sub> atmosphere.

#### Example 2: Immunoblotting procedures

Subconfluent cells in 10-cm dishes cells were washed twice with cold PBS and solubilized with PI/RIPA (1% Triton X-100, 1% deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate-pH 7.4, 1% Aprotinin, 1 mM phenylmethanolsulfonyl fluoride, 2 mM EDTA, 10 mM sodium pyrophosphate, 400 mM sodium orthovanadate, and 10 mM iodoacetamide buffer. Proteins were separated by proper concentration of SDS-PAGE and transferred to nitrocellulose 15 membranes (Nitrobind, MSI). Membranes were incubated overnight with the blocking buffer (0.5% non-fat milk and 5% goat serum in PBS). For immunoblotting (Western blot) analysis, antibodies were diluted in PBS containing 0.1% non-fat milk and 1% goat serum. All polyclonal sera and 20 secondary HRP-conjugated antibodies (Boehringer Mannheim) were used at a 1:5000 dilution. Bands were visualized using the ECL assay (Amersham).

#### Example 3: Expression and Purification of His-tagged 7.16.4

LB media (150 ml) containing 50 µg/ml ampicillin was 25 inoculated with a 15 ml overnight culture of *E. coli* DH5 α and maintained at 30°C until an optical density of 0.5 (600 nm) was obtained. Isopropyl-β-(D-thiogalactopyranoside) (IPTG) was added at a final concentration of 1 mM to induce His-tagged 7.16.4ScFv. After 3 hours, cells were 30 harvested and lysed by freezing and thawing and resuspended in 10 ml of urea lysis buffer (10 mM Tris-Cl, pH 8.0, 0.1 M NaH<sub>2</sub>PO<sub>4</sub>, 1 M NaCl, 8 M urea) supplemented with 0.5 mM phenylmethanolsulfonyl fluoride, 2 µM pepstatin A, and 2 µM

leupeptin. The insoluble cellular debris was removed by centrifugation (12,000 x g for 15 minutes, followed by 12,000 x g for 30 minutes). The clarified supernatant was mixed with 2 ml of Ni-NTA agarose followed by gentle shaking on ice for 1 hour. The mixture was loaded into an empty column and unbound protein was washed with urea lysis buffer (pH 6.3) at 4°C. His-tagged protein was eluted with urea lysis buffer (pH 4.5), and eluate fractions were examined by SDS-PAGE and FACS analysis as described in Example 4. Proteins in the peak fractions were pooled and dialyzed against TKC buffer (50 mM Tris-Cl, pH 8.0, 100 mM KCl, 10 mM CaCl<sub>2</sub>, 1 mM EDTA, 0.1 mM PMSF).

#### **Example 4: Confirmation of 7.16.4ScFv binding by FACS**

##### **Analysis**

Cell-surface antigens were detected by Fluorescence-activated cell sorting (FACS). B104-1-1 cells (about 5x10<sup>5</sup>) were incubated (30 minutes at 4°C) in 200 µl of FACS buffer (PBS containing 0.5% bovine serum albumin and 0.02% sodium azide) containing the purified 7.16.4ScFv. Cells were then washed in FACS buffer and incubated (30 minutes, 4°C) with anti-His antibody (Invitrogen). After this second incubation, cells were washed again in FACS buffer and further incubated with FITC-labeled IgG against mouse immunoglobulins (Sigma Chemical Co., St. Louis, MO). After washing with FACS buffer, the cell pellet was suspended in PBS buffer and processed for analysis by a FACS scan flow cytometer (Becton Dickinson). For each sample, 10,000 viable cells were gated following size (forward scatter, FSC) and granularity (side scatter, SSC) parameters and analyzed with CellQuest Software (Becton Dickinson). For competitive binding, B104-1-1 cells were first incubated with the monoclonal antibody 7.16.4 in the presence of different concentrations of 7.16.4ScFv. Cells were then washed in FACS buffer and further incubated with

FITC-labeled IgG against mouse immunoglobulins (Sigma Chemical Co.) before analysis on the flow cytometer.

**Example 5: Attachment of ds-cDNA to Antibody or Fv or CDR**

5 A cdna oligonucleotide of the following sequence  
(GAGAGAGGATCCAAGTACTAATACGACTCACTATAGGGCCGAGAGCCGAGAAGAAAGA  
CGTTTTTTTTTTT (SEQ ID NO:1)) was designed with an  
activatable amine at the 5' end to allow for covalent  
coupling to primary amines and a T7 promotor site  
10 (TAATACGACTCACTATAGGG (SEQ ID NO:2)) used to direct the  
synthesis of RNA from the cdna template through the  
enzymatic activity of T7 RNA polymerase. For attachment of  
1 µg of antibody to 30 µg of ds cdna, or 1 µg CDR to 0.1 µg  
ds cdna or 1 µg ScFv to 3 µg ds cdna, an equal volume of  
15 0.1% glutaraldehyde was added in 10 µl aliquots. The  
solution was mixed on a rotation device for 3 hours at room  
temperature. Ethanolamine (1 M; 1/20 volume; pH 7.5) was  
then added. The solution was mixed for an additional 2  
hours at room temperature. The protein-DNA complex was  
20 stored at 4°C.

**Example 6: RNA amplification**

RNA amplification was performed in 96-well plates  
containing the first Ab-Antigen complex. The following  
reagents were added: 1X RNA amplification buffer (40 mM  
25 Tris-base, pH 7.5, 7 mM MgCl<sub>2</sub>, 10 mM NaCl, 2 mM Spermidine;  
5 mM DTT; 250 µM ATP, UTP, GTP; 5 µM CTP; 0.5 µl RNAsin;  
1000 U T7 RNA Polymerase and 3 µl P<sup>32</sup>-CTP. The solution was  
then incubated for 4 hours at 37°C. The RNA product was  
removed from the wells and 3 µl was added to 3 µl of  
30 reaction stop solution (95% formamide, 10 mM EDTA, 0.1%  
bromophenol blue, 0.1% xylene cyanol) and electrophoresed  
on a 15% denaturing polyacrylamide gel. The gel was  
apposed to a phosphoimager screen for 5-60 minutes and



developed on a phosphoimager at 100 uM resolution.

**Example 7: Standard Sandwich ELISA as a Control Method**

Ninety-six well microtiter plates (Nunc-Immuno Plate, MaxiSorp™) were coated with the 1E1 (a monoclonal antibody directed against the extracellular domain of human p185) by incubating plates overnight at 4°C with 100 µl of coating buffer (antibody concentration: 5 µg/ml). Plates were then washed three times with PBS containing 0.2% TWEEN 20 (PBS-T; 200 µl/well), blocked with PBS containing 0.5% FBS and 0.2% TWEEN 20 (200 µl/well) for 1 hour at room temperature, and washed again three times with PBS-T (200 µl/well). In parallel, 100 µl of a serial dilution of T6-17 cell lysates were added to the plate and incubated for 2 hours at room temperature. After this incubation step, plates were washed with PBS-T (six times, 200 µl/well) and incubated with humanized anti-p185Her2 antibody 4D5 (150 ng/ml, 50 µl/well) for 2 hours at room temperature. Subsequently, plates were washed six times and incubated with 50 µl of anti-human IgG-HRP (Zymed; final dilution: 1:10,000) for another 2 hours at room temperature. Enzymatic reactions were carried out at room temperature by adding TMB (2.5 mM each in 0.1 M phosphate-citrate buffer, pH 5.0) (100 µl of each reagent/well). Reactions were stopped after 15-60 minutes by the addition of 50 µl of 1 M H<sub>2</sub>SO<sub>4</sub>. Color development was measured at 450 nm using the ELISA reader.

What is Claimed is:

1. A method for detecting molecules expressing a selected epitope in a sample comprising:

- (a) immobilizing a molecule expressing a selected epitope in a sample to a selected surface;
- (b) contacting the surface with an epitope detector so that the epitope detector binds to immobilized molecules on the surface; and
- (c) detecting any epitope detector bound to the surface, wherein bound epitope detector is indicative of molecules expressing the selected epitope in the sample.

2. The method of claim 1 wherein the molecule expressing a selected epitope is immobilized to the selected surface via binding to an epitope anchor on the surface which is specific for the selected epitope.

3. The method of claim 1 wherein the epitope detector comprises a universal epitope detector which detects a general epitope.

4. The method of claim 1 wherein the detected molecule is post-translationally modified.

5. A system for the detection of molecules expressing a selected epitope comprising:

- (a) a selected surface on which a molecule expressing a selected epitope is or can be immobilized; and
- (b) an epitope detector comprising a single chain Fv for the selected epitope or a constrained epitope specific CDR either of which have been modified to allow for attachment of oligonucleotides.

6. The system of claim 5 further comprising an epitope anchor for immobilizing the molecule to the

selected surface, said epitope anchor being specific for the selected epitope.

7. The system of claim 5 wherein the epitope detector comprises a universal epitope detector which  
5 detects a general epitope.

8. A kit for the detection of molecules expressing a selected epitope comprising an epitope detector comprising a single chain Fv for the selected epitope or a constrained epitope specific CDR.

10 9. The kit of claim 8 further comprising an epitope anchor specific for the selected epitope.

10 10. The kit of claim 8 wherein the single chain Fv or the constrained epitope specific CDR have been modified  
15 for attachment of oligonucleotides.

11. The kit of claim 8 wherein the epitope detector comprises a universal epitope detector which detects a general epitope.

005220 94042500

## ABSTRACT

Methods, systems and kits are provided for detecting molecules expressing a selected epitope in a sample through use of an epitope detector containing a single chain Fv for the selected epitope or a constrained epitope specific CDR attached to an oligonucleotide.

Docket No.  
PENN-0708

# Declaration and Power of Attorney For Patent Application

## English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**A RAPID, SENSITIVE AND QUANTITATIVE METHOD FOR IMMUNO-DETECTION OF EPITOPES ON MOLECULES USING A SINGLE CHAIN FV FOR THE EPITOPE OR A CONSTRAINED EPITOPE**

the specification of which

(check one)

☒ is attached hereto.

☐ was filed on \_\_\_\_\_ as United States Application No. or PCT International Application Number \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

I hereby claim the benefit under 35 U. S. C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, C. F. R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

\_\_\_\_\_  
(Application Serial No.)

\_\_\_\_\_  
(Filing Date)

\_\_\_\_\_  
(Status)  
(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*

**Jane Massey Licata, Reg.No. 32,257**

**Kathleen A. Tyrrell, Reg. No. 38,350**

**Laura Plunkett, Reg. No. 45,015**

**of the firm**

**Law Offices of Jane Massey Licata  
66 E. Main Street  
Marlton, NJ 08053**

Send Correspondence to: **Jane Massey Licata  
Law Offices of Jane Massey Licata  
66 E. Main Street  
Marlton, NJ 08053**

Direct Telephone Calls to: *(name and telephone number)*  
**Jane Massey Licata Tel: 856-810-1515**

Full name of sole or first inventor <b>Mark I. Greene</b>	
Sole or first inventor's signature	Date
Residence <b>Penn Valley, PA</b>	
Citizenship <b>US</b>	
Post Office Address <b>300 Righters Mill Road</b>	
<b>Penn Valley, PA 19071</b>	

Full name of second inventor, if any <b>James Eberwine</b>	
Second inventor's signature	Date
Residence <b>Philadelphia, PA</b>	
Citizenship <b>US</b>	
Post Office Address <b>3918 Henry Avenue</b>	
<b>Philadelphai, PA 19129</b>	

Full name of third inventor, if any <b>Janet Estee Kacharmina</b>	
Third inventor's signature	Date
Residence <b>Philadelphia, PA</b>	
Citizenship <b>US</b>	
Post Office Address <b>7312 Bryan Street</b>	
<b>Philadelphia, PA 19119</b>	

Full name of fourth inventor, if any <b>Hong Tao Zhang</b>	
Fourth inventor's signature	Date
Residence <b>Devon, PA</b>	
Citizenship <b>China</b>	
Post Office Address <b>377 Poplar Avenue, #N488</b>	
<b>Devon, PA 19333</b>	

Full name of fifth inventor, if any	
Fifth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	

Full name of sixth inventor, if any	
Sixth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	



# SEQUENCE LISTING

<110> Greene, Mark I.

Eberwine, James

Kacharina, Janet Estee

Zhang, Hong Tao

<120> A Rapid, Sensitive and Quantitative Method for  
Immuno-Detection of Epitopes on Molecules using a  
Single Chain Fv for the Epitope or a Constrained  
Epitope Specific CDR

<130> PENN-0708

<140>

<141>

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 70

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

<400> 1

gagagaggat ccaagtacta atacgactca ctatagggcc gagagccgag aagaaagacg 60  
tttttttttt 70

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>


<223> Description of Artificial Sequence: Synthetic

<400> 2

taatacgact cactataggg

20

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



SCAN 8

Q: Application deficiencies were found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present:  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present:  
for scanning. (Document title)

☒ Scanned copy is best available. Cf verified statement.